#### Water Quality Program


Figures for Oregon Forest Industry Council's Questions/Concerns Re: Protecting Cold Water Criterion


June 19<sup>th</sup>, 2014

Oregon Department of Environmental Quality



## Figure 1: Energy Fluxes @ RipStream Site 5556






Modeled heat energy fluxes **pre harvest** at stream km 1.25 in the treatment reach on July 31, 2003.

Modeled heat energy fluxes **post harvest** at stream km 1.25 in the treatment reach on July 31, 2003.

Oregon Department of Environmental Quality



# Figure 2: Field Measured Energy Fluxes



Thermal energy fluxes at noon on July 20, 1997 on a <u>fully</u> <u>shaded</u> study stream in the H.J. Andrews Experimental Forest. Thermal energy lost is -149 W\*m<sup>-2</sup>.


Thermal energy fluxes at noon on July 20, 1997 on an **unshaded** study stream in the H.J. Andrews Experimental Forest. **Thermal energy gained is 580 W\*m**-2.

Johnson 200

Oregon Department of Environmental Quality



# Figure 3: Heat Source Results for Argue Creek




Simulated change in longitudinal 7-day average daily maximum (7DADM) temperatures from harvest at RipStream site 7854, holding all factors constant except vegetation.

Oregon Department of Environmental Quality



### Figure 4: Heat Source Results for Drift Creek Trib




Simulated change in longitudinal net energy fluxes from harvest at RipStream site 5556, holding all factors constant except vegetation.

Results only include effect of the Harvest Unit 1 in RipStream study area.

Oregon Department of Environmental Quality



### Figure 5: Heat Source Results for Drift Creek Trib




Simulated change in longitudinal 7-day average daily maximum (7DADM) temperatures from harvest at RipStream site 5556, holding all factors constant except vegetation.

Results only include effect of the Harvest Unit 1 in RipStream study area.

Oregon Department of Environmental Quality



### Figure 6: Heat Source Results for Drift Creek Trib




Simulated change in longitudinal 7-day average daily maximum (7DADM) temperatures from harvest at RipStream site 5556, holding all factors constant except vegetation.


Results include effects from the harvest unit in RipStream study area (Harvest Unit 1) and a second harvest unit downstream of the study area (Harvest Unit 2).

Oregon Department of Environmental Quality



#### Figure 7: Change Detection Analysis





Kennedy et al. 2010.

Percent area disturbed in 2006 = 17%

Includes any harvest 10 years prior from 1996-2006.

Color = harvested 1996 - 2006

White = Non forested (Ag/Rural residential, meadow)

Dark Green = not harvested 1985 - 2006.

Light Green = harvested 1985-1996. (Not counted in

total

DEQ


J. U

### Figure 8: Thermal Recovery in Middle Siletz River

Percent of riparian area in thermal recovery (≤10 years post-disturbance) as a percentage of the land use in question.

Percentages in the legend are the amount of the total watershed area in that land use.

Riparian areas are land within 100ft of streams.




Oregon Department of Environmental Quality



### Figure 9: Thermal Recovery in Middle Siletz River

Percent of riparian area in thermal recovery (≤10 years post-disturbance) as a percentage of the total watershed riparian area.

Riparian areas are land within 100ft of streams.



Oregon Department of Environmental Quality




### Figure 10: Thermal Recovery in Drift Creek

Percent of riparian area in thermal recovery (≤10 years post-disturbance) as a percentage of the land use in question.

Percentages in the legend are the amount of the total watershed area in that land use.

Riparian areas are land within 100ft of streams.




Oregon Department of Environmental Quality



### Figure 11: Thermal Recovery in Drift Creek

Percent of riparian area in thermal recovery (≤10 years post-disturbance) as a percentage of the total watershed riparian area.

Riparian areas are land within 100ft of streams.



Oregon Department of Environmental Quality




### Figure 12: Thermal Recovery in Lake Creek

Percent of riparian area in thermal recovery (≤10 years post-disturbance) as a percentage of the land use in question.

Percentages in the legend are the amount of the total watershed area in that land use.

Riparian areas are land within 100ft of streams.




Oregon Department of Environmental Quality



### Figure 13: Thermal Recovery in Lake Creek

Percent of riparian area in thermal recovery (≤10 years post-disturbance) as a percentage of the total watershed riparian area.

Riparian areas are land within 100ft of streams.



Oregon Department of Environmental Quality

